

ORIGINAL ARTICLE

Invasive pulmonary aspergillosis in patients with influenza infection: A retrospective study and review of the literature

Linna Huang¹  | Nannan Zhang^{1,2} | Xu Huang¹ | Shuyu Xiong¹ |
Yingying Feng¹ | Yi Zhang¹ | Min Li¹ | Qingyuan Zhan¹

¹Department of Pulmonary and Critical Care Medicine, Centre for Respiratory Diseases, China-Japan Friendship Hospital, Beijing, P.R. China

²Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, P.R. China

Correspondence

Qingyuan Zhan, Department of Pulmonary and Critical Care Medicine, Centre for Respiratory Diseases, China-Japan Friendship Hospital, No. 2 Yinghua East Road, Chaoyang District, Beijing 100029, P.R. China.
Email: drzhanqy@163.com

Funding information

Dr Zhan was supported by a grant from the National Key Research and Development Programme—Major Chronic Non-Communicable Diseases' Prevention and Control (QML 2016YFC1304300) and the Beijing Municipal Science and Technology Project (Z16100000516116).

Abstract

Introduction: There has been a rapid increase in the number of influenza and invasive pulmonary aspergillosis (IPA) co-infection.

Objectives: To explore the risk factors and predictors of a poor prognosis in influenza and IPA co-infection.

Methods: We included patients with confirmed influenza during the 2017-2018 influenza season and cases of influenza and IPA co-infection in the literature.

Results: A total of 64 patients with influenza infection were admitted to ICU. Of these patients, 18 were co-infected with IPA. Others were assigned to the control group ($n = 46$). A total of 45 patients from the literature were added to the IPA group ($n = 63$). A multivariate logistic regression suggested that influenza patients who were given steroids after ICU admission, who had a white blood count (WBC) of more than $10 \times 10^9/L$ on ICU admission and whose CT findings manifested as multiple nodules and cavities might have a higher risk of developing IPA. Compared to survivors, non-survivors had higher sequential organ failure assessment (SOFA) scores (16 ± 4 points vs 8 ± 4 points, $P < 0.001$), lower $CD4^+$ T cell counts on ICU admission [315 (83 - 466) cells/ μL vs 152 (50 - 220) cells/ μL , $P = 0.031$] and more requirement extracorporeal membrane oxygenation (ECMO) support [13 (50%) vs 7 (18.9%), $P = 0.015$].

Conclusions: Influenza patients who are given steroids after ICU admission, who have WBCs of greater than $10 \times 10^9/L$ on ICU admission, and whose CT imaging shows multiple nodules and cavities might have a high risk of IPA. Higher SOFA scores, $CD4^+$ T cell counts lower than 200 cells/ μL on ICU admission and more ECMO requirement might be predictors of a poor prognosis.

KEYWORDS

clinical presentations, influenza, invasive pulmonary aspergillosis (IPA), prognosis, risk factor

Abbreviations: AKI, acute kidney injury; ARDS, acute respiratory distress syndrome; CHF, chronic heart failure; CRBSI, catheter-related bloodstream infection; CT, computed tomography; CTD, connective tissue disease; DM, diabetes mellitus; ECMO, extracorporeal membrane oxygenation; GGO, ground glass opacity; GM, galactomannan; IPA, invasive pulmonary aspergillosis; IPPV, invasive positive pressure ventilation; LRT, lower respiratory tract; MICU, medical intensive care unit; PCR, polymerase chain reaction; PCT, procalcitonin; SOFA, sequential organ failure assessment; VAP, ventilation associated pneumonia; WBC, white blood count.

Linna Huang and Nannan Zhang contributed equally to the article.

1 | INTRODUCTION

Co-infection as a common complication of influenza infection that has been well described, and it often contributes to increased morbidity and mortality.¹ Bacterial pathogens, often including *Streptococcus* and *Staphylococcus aureus*, have become considerable issues,² while co-infection with *Aspergillus* has typically been ignored. However, it might play a role in severe influenza cases.

Invasive *Aspergillus* infections are a well-known burden for immunocompromised patients. While *Aspergillus* is not traditionally thought of as a pathogen capable of invasive infection following a viral respiratory infection, since 2010, there has been a rapid increase in the amount of research investigating influenza and subsequent invasive *Aspergillus* co-infection.³

Influenza may represent a novel host risk factor for this invasive fungal infection. Meanwhile, the mortality rate among co-infected patients has been found to be approximately 50%-60%,⁴ which is approximately 5 times that of hospitalized patients with influenza alone.

Invasive pulmonary aspergillosis (IPA) co-infections with influenza have been sporadically reported in case reports and small case series,⁵⁻⁷ but no systemic research has yet been aimed at investigating the clinical characteristics of IPA co-infections with influenza, how to recognize it earlier and its poor prognostic factors.

Our study included patients with IPA complicating severe influenza infections who were admitted to the medical intensive care unit (MICU) of the China-Japan Friendship Hospital during the 2017-2018 influenza season. In addition, cases from a comprehensive review of the English literature were added to provide a better and more comprehensive view of the clinical features, potential risk factors and predictors of the prognosis of influenza combined with IPA.

2 | MATERIALS AND METHODS

2.1 | Study population and group divisions

2.1.1 | Our cases

We retrospectively included all confirmed influenza-infected patients with respiratory failure admitted to the MICU of the China-Japan Friendship Hospital from November 2017 to March 2018.

2.1.2 | Cases from the literature

In addition to our current cases, we searched the English language published literature using PubMed/Medline with the search terms 'influenza' and 'aspergillus' or 'aspergillosis' from January 2009 to March 2018.^{1,5-18}

2.1.3 | Group division

We divided the patients into two groups. The case group included patients with confirmed influenza infection who subsequently became infected with proven and/or probable IPA, and the control group included patients with confirmed influenza infection who showed no evidence of *Aspergillus* infection while hospitalized.

Cases involving nonspecific viral infections or without confirmatory influenza testing, reports with insufficient patient information and with possible IPA or *Aspergillus* colonization were excluded.

2.2 | Diagnostic criteria

2.2.1 | Definition of influenza virus infection

All patients with influenza had testing performed using a nasopharyngeal swab and lower respiratory tract (LRT) specimens. Three methods were used for a laboratory diagnosis, namely a polymerase chain reaction or a respiratory viral culture along with serological testing.^{3,19}

2.2.2 | Definition of IPA

Proven IPA is defined as microscopic evidence of dichotomous branching hyphae with a positive culture for *Aspergillus* spp through an endobronchial biopsy, irrespective of host factors or clinical features.²⁰ Probable IPA requires a host factor, clinical features and mycological evidence of aspergillosis. One of the three criteria²⁰⁻²² for an IPA diagnosis is host factors; however, these criteria were largely created for immunosuppressed hosts, and influenza-related aspergillosis may occur in previously normal hosts. Thus, the host factors criterion was not required in our study. The clinical signs and symptoms, radiological findings and mycological data should be met according to our newly proposed 'Modified Bulpa Criteria'²³ (Additional File 1).

2.3 | Data collection and analysis

Data were retrospectively collected from electronic medical records, including demographics, underlying diseases and the use of immunosuppressive agents and steroids before admission. Data on the use of steroids during the hospitalizations were recorded. Laboratory examinations included the influenza-type, white blood count (WBC) and lymphocyte and CD4⁺ T lymphocyte counts. All respiratory culture data from the hospitalizations were evaluated. Radiological examinations, mainly chest computed tomography (CT), were reviewed. Complications during hospitalizations and needs for organ support were also recorded. Data regarding lengths

TABLE 1 Characteristics, radiological findings and mycological data of our patients

Case	Patients characteristics				Time from influenza to IPA	Grade of IPA diagnosis	Mycological data				
	Age	Sex	Underlying diseases	IPA			Radiological findings	Culture	BALF GM	Serum GM	Histo evidence
1	87	F	Lung disease	21	Probable	Tree in bud	<i>Aspergillus fumigatus</i>	+	-	Not done	Improved
2	61	M	CTD	27	Probable	Nodules, mass, patchiness	None	+	-	Not done	Died
3	81	F	None	13	Probable	Tree in bud, GGO	None	-	+	Not done	Improved
4	51	M	None	6	Probable	Nodules, patchiness, large consolidation, cavities, reversed halo sign	None	+	-	Not done	Improved
5	49	M	None	22	Probable	Large consolidation, GGO	None	+	-	Not done	Died
6	46	M	DM	9	Probable	Nodules, patchiness, large consolidation, cavity	<i>Aspergillus fumigatus</i>	+	+	Not done	Improved
7	41	F	None	8	Probable	Nodules, patchiness, large consolidation, halo sign, cavity, GGO	<i>Aspergillus flavus</i>	+	-	Not done	Improved
8	51	M	DM	9	Probable	Patchiness, large consolidation, air-crescent sign, GGO	<i>Aspergillus fumigatus</i>	+	+	Not done	Improved
9	51	M	DM	18	Probable	Patchiness, large consolidation, cavity	None	+	+	Not done	Improved
10	63	M	CTD	16	Probable	Patchiness	<i>Aspergillus fumigatus</i>	-	+	Not done	Died
11	16	M	None	13	Probable	Nodules, patchiness, large consolidation, reversed halo sign, cavity	<i>Aspergillus fumigatus</i>	+	+	Not done	Died
12	46	F	DM	15	Proven	Nodules, patchiness, large consolidation, halo sign	<i>Aspergillus fumigatus</i> ; <i>Aspergillus flavus</i>	+	+	Autopsy	Died
13	65	F	DM, lung diseases	7	Probable	Nodules, cavities	<i>Aspergillus fumigatus</i>	+	-	Not done	Improved
14	66	M	DM	23	Probable	Nodules, patchiness, cavities	None	+	+	Not done	Died
15	35	M	DM	10	Probable	Nodules, patchiness, large consolidation, cavities	<i>Aspergillus fumigatus</i>	+	-	Not done	Died
16	52	M	None	12	Probable	Nodules, patchiness, halo sign	<i>Aspergillus flavus</i>	+	+	Not done	Improved
17	76	F	DM, CHF, lung disease	2	Probable	Nodules, patchiness	None	+	-	Not done	Improved
18	75	M	Malignancy	5	Probable	Patchiness, large consolidation, cavity	None	+	-	Not done	Died

Abbreviations: CHF, chronic heart failure; CTD, connective tissue disease; DM, diabetes mellitus; GGO, ground glass opacity; GM, galactomannan; F, female; IPA, invasive pulmonary aspergillosis; M, male.




FIGURE 1 Flowchart. A flowchart illustrating the enrolment of patients in our study. From November 2017 to March 2018, 64 critically ill patients with confirmed influenza were admitted to the MICU at the China-Japan Friendship Hospital due to respiratory failure. Of these patients, 18 suffered from IPA, and the other 46 were designated as the control group. We added the IPA cases from the literature (45 cases) to our IPA cases (18 cases) to create the case group ($n = 63$). Out of the 63 patients included in the case group, 37 survived and 26 died

of MICU stays, total hospitalization days and mortality before discharge were collected as well. For those with aspergillosis, data on the number of days between the diagnoses of influenza and aspergillosis, the *Aspergillus* species from the LRT cultures, serum and/or BALF galactomannan levels and antifungal therapy information were recorded.

We first described the overview of this influenza pandemic (from November 2017 to March 2018) and IPA coinfection. We then added the IPA cases from the literature to our IPA cases to create the case group, while the other patients from our cases were placed in the control group. We compared the two groups in terms of demographics, underlying diseases, use of immunosuppressive agents and steroids before admission, steroid use during hospitalization and laboratory and radiological data to investigate the clinical presentation of influenza combined with IPA. Then, a logistic regression analysis was performed to explore the condition's risk factors. Finally, we compared survivors and non-survivors of IPA to determine predictors of a poor prognosis.

2.4 | Statistical analysis

All of the analyses were carried out using SPSS 16.0 software. Normally distributed continuous variables were expressed as $\bar{x} \pm SD$ and were compared using *t*-tests. Non-normally distributed continuous variables were expressed as medians and quartiles and were compared using Wilcoxon rank-sum tests. Categorical variables were compared using χ^2 tests. A multivariate logistic regression analysis was used to identify independent risk factors and

poor prognosis for influenza combined with IPA. Variables in bivariate analyses with *P* value less than 0.2, as well as the variables thought to be meaningful in practice even though with no statistical significance are all included in the model. Power analysis was performed by PASS 16 to verify whether the sample size was adequate to identify risk factors and poor prognosis in multivariate logistic regression analysis.

3 | RESULTS

3.1 | Overview of patients with influenza in the MICU in the 2017 to 2018 influenza season

From November 2017 to March 2018, 64 critically ill patients with confirmed influenza were admitted to the MICU at the China-Japan Friendship Hospital due to respiratory failure; the mortality rate for all of these influenza cases was 43.8%.

Among these patients, 18 suffered from IPA, giving an incidence of 28.1%. One patient was diagnosed with proven IPA using a trans-bronchial lung biopsy. The other 17 patients were diagnosed with probable IPA, and the mean time from the influenza diagnosis to the IPA diagnosis was 13 (± 7) days. The average age of the patients was 52 (± 18) years, 66.7% were male and the mean sequential organ failure assessment (SOFA) score was 8 (± 4) points. Ten patients (55.6%) had underlying diseases, one (5.6%) had a history of immunosuppressant use and eight (44.4%) reported steroid use in the last month. Steroids were used after MICU admission in five patients. Invasive positive pressure ventilation (IPPV) was required for 16 patients (88.9%), and the average duration of its use was 14 (5-33) days. Rescue ventilation strategies and extracorporeal membrane oxygenation (ECMO) were needed for 10 (55.7%) and 4 (22.2%) patients, respectively; the average duration of ECMO use was 10 (6-23) days. The mean lengths of ICU and total hospital stays were 16 (6-35) days and 19 (10-36) days, respectively. The mortality rate was 44.4% (Table 1).

3.2 | Comparisons between IPA and non-IPA patients

We added the IPA cases from the literature (45 cases) to our IPA cases (18 cases) to create the IPA group ($n = 63$), while the other patients from our set of cases ($n = 46$) were designated as the control group (Figure 1).

3.2.1 | Patient characteristics and outcomes between the two groups

General information

The IPA and control groups did not differ significantly in terms of the number of cases of obesity, diabetes, underlying

lung disease, chronic heart failure, chronic kidney disease, steroid use and accumulated steroid dosage prior to ICU admission. Compared to the control group, the IPA group included more male patients (68.3% vs 47.8%, $P = 0.032$), was older (63 ± 13 years vs 57 ± 18 , $P = 0.044$) and had more severe conditions (SOFA scores: 10 ± 3 points vs 7 ± 4 points, $P < 0.001$). The numbers of patients with a malignancy (23.8% vs 6.5%, $P = 0.016$), with immunosuppressant use in the past month (46% vs 19.6%, $P = 0.005$), and who were given steroids after ICU admission [234 (0-400) mg vs 89 (0-120) mg, $P = 0.036$] were higher in the IPA group than in the non-IPA group. Moreover, compared to the control group, the accumulated dose of steroids was much higher in the IPA group (Table 2).

Laboratory examinations and radiological findings

The IPA group had a higher WBC [$(10.2 \pm 4.6) \times 10^9/L$ vs $(6.1 \pm 3.3) \times 10^9/L$, $P < 0.001$], and lower lymphocyte count [$(0.47 \pm 0.52) \times 10^9/L$ vs $(0.67 \pm 0.46) \times 10^9/L$, $P = 0.035$] on ICU admission. CD4⁺ T cell counts on ICU admission and types of influenza did not differ significantly between the two groups. Patchiness was seen more often in the CT imaging of non-IPA patients, while nodules (51.9% vs 21.1%, $P = 0.003$), and especially multiple nodules distributed along the airway bundles and cavities (30.8% vs 10.5%, $P = 0.013$) were seen more often in IPA patients. The incidences of masses, large consolidations and halo signs were similar between the two groups (Table 2).

Clinical outcomes

Despite there being no significant differences in mortality between the two groups, the lengths of ICU [25 (13-39) vs 12 (7-20), $P = 0.005$] and total hospital stays [28 (14-58) vs 18 (10-27), $P = 0.002$] was higher in the IPA group than in the control group (Table 2).

3.2.2 | Organ support needs and complications between the two groups

Compared to the control group, more patients in the IPA group needed rescue ventilation strategies (42.9% vs 21.7%, $P = 0.021$), renal replacement therapies (49.2% vs 28.3%, $P = 0.033$), and high-dose vasoactive drugs (68.3% vs 41.3%, $P = 0.005$). The numbers of patients needing IPPV and ECMO support did not differ significantly between the two groups. The duration of IPPV was significantly longer in the IPA group than in non-IPA group [22 (9-27) days vs 17 (7-18) days, $P = 0.047$], while the durations of renal replacement therapy and ECMO support did not differ significantly. For complications, patients in the IPA group were more prone to developing septic shock (with incidences of 66.7% vs 41.3%, $P < 0.001$). Other complications, such as the incidences of ventilator-associated pneumonia (VAP),

catheter-related bloodstream infection (CRBSI), bacteremia, barotrauma and haemorrhage were similar between the two groups (Table 3).

3.3 | Subgroup analysis: comparison between survivors and non-survivors in the IPA group

Out of the 63 IPA patients included in this study, 26 died, giving a mortality rate of 41.3%. Compared to survivors, non-survivors were older (60 ± 15 years vs 67 ± 9 years, $P = 0.040$), had higher SOFA scores (16 ± 4 points vs 8 ± 4 points, $P < 0.001$) and had lower CD4⁺ T cell counts on ICU admission [315 (83-466) cells/ μ L vs 152 (50-220) cells/ μ L, $P = 0.031$]. All of the non-survivors had influenza type A (100% vs 70.3%, $P = 0.002$), and more non-survivors required IPPV (96.2% vs 54.1%, $P < 0.001$), rescue ventilation strategies (57.7% vs 32.4%, $P = 0.046$) and ECMO support (50% vs 18.9%, $P = 0.015$) compared to survivors in the IPA group (Table 4).

3.4 | Multivariate logistic regression analysis for independent risk factors and the predictors of a poor prognosis for IPA in patients with influenza

To investigate the roles of potential confounding risk factors and predictors of poor prognosis of IPA, a multivariate logistic regression analysis was performed. We found that influenza patients who were given steroids after ICU admission, who had a WBC of more than $10 \times 10^9/L$, and whose CT imaging showed multiple nodules and cavities might have a higher incidence of IPA. Higher SOFA scores, CD4⁺ T cell counts lower than 200 cells/ μ L on ICU admission and more ECMO requirement might be predictors of a poor prognosis (Table 5).

The systemic steroid application was commonly regarded as the risk factor for influenza and IPA infections in studies before.⁸ In this way, the 'Steroids after ICU admission' was identified as primary factor for power analysis. The power was 0.77, which might verify the sample size was adequate to identify risk factors.

4 | DISCUSSION

The biggest strength of our study is the large group of literature-based cases of IPA following influenza infection that was included. Additionally, this is the first study to systematically and comprehensively provide an overview of the clinical presentation of influenza combined with IPA and to explore the possible risk factors for its diagnosis and predictors of a poor prognosis.

TABLE 2 Patient characteristics and outcomes between the two groups

	IPA (n = 63)	Non-IPA (n = 46)	P Value
<i>General information</i>			
Age, years, mean ± SD	63 ± 13	57 ± 18	0.044*
Sex (men), number (%)	43 (68.3)	22 (47.8)	0.032*
BMI > 30 kg/m ² , number (%)	5 (7.9)	5 (10.9)	0.600
SOFA, mean ± SD	10 ± 3	7 ± 4	0.001#
Diabetes, number (%)	19 (30.2)	12 (26.1)	0.642
Underlying lung disease, number (%)	13 (20.6)	5 (10.9)	0.175
Chronic heart failure, number (%)	9 (14.3)	3(6.5)	0.201
Chronic kidney disease, number (%)	2 (3.2)	2 (4.3)	0.748
Malignancy, number (%)	15 (23.8)	3 (6.5)	0.016*
Immunosuppressant use in past month, number (%)	15 (23.8)	3 (6.5)	0.016*
<i>Systemic steroid use</i>			
Before ICU admission, number (%)	23 (36.5)	13 (28.3)	0.366
Accumulated dosage before ICU, mg, median (IQR)	242 (0-300)	238 (0-1092)	0.977
After ICU admission, number (%)	29 (46.0)	9 (19.6)	0.005*
Accumulated dosage after ICU, mg, median (IQR)	234 (40-400)	89 (0-120)	0.036*
<i>Laboratory examination</i>			
WBCs on ICU admission, *10 ⁹ /L, mean ± SD	10.2 ± 4.6	6.1 ± 3.3	<0.001#
Lymphocytes on ICU admission, *10 ⁹ /L, mean ± SD	0.47 ± 0.52	0.67 ± 0.46	0.035*
CD4 ⁺ T cells on ICU admission, cell/μL, mean ± SD	239 (82-416)	320 (209-458)	0.373
Influenza A, number (%)	52 (82.5)	36 (78.3)	0.699
Influenza B, number (%)	11 (17.5)	11 (23.9)	0.384
<i>Radiological findings</i>			
Patchiness, number (%)	34 (65.4)	37 (97.4)	<0.001#
Nodules, number (%)	27 (51.9)	8 (21.1)	0.003*
Mass, number (%)	1 (1.9)	1 (2.6)	0.811
Large consolidations, number (%)	24 (46.2)	11 (28.9)	0.098
Halo sign, number (%)	5 (9.6)	0 (0.0)	0.051
Cavity/air-crescent sign, number (%)	16 (30.8)	4 (10.5)	0.013*
<i>Outcomes</i>			
Length of ICU stay, days (IQR)	25 (13-39)	12 (7-20)	0.005*
Length of hospitalization, days (IQR)	28 (14-58)	18 (10-27)	0.002*
Mortality, number (%)	26 (41.3)	20 (43.4)	0.818

Abbreviations: BMI, body mass index; ICU, intensive care unit; IQR, interquartile range; SD, standard deviation; SOFA, sequential organ failure assessment; WBC, white blood count.

The bold values indicates P values from 0.01 to 0.05.

Note. The steroid doses were converted to prednisone doses.

*P < 0.05;

#P < 0.01.

Our study revealed that the incidence of IPA co-infection in the influenza season examined here was 28.1%. A study by Nancy F. in 2016³ showed that 5 (62.5%) of 8 ICU influenza cases were classified as an invasive disease, which was much higher than in our cases. A rising number of recent cases have revealed that influenza might represent a novel host risk factor for invasive fungal infection. The pathogenesis of IPA in the setting of influenza infection may occur as follows. The

influenza virus may cause severe and diffuse damage to the respiratory mucosa and disrupt normal ciliary clearance,^{24,25} allowing for fungal invasion. Influenza may also impair local phagocytosis by alveolar macrophages as well as reduce natural killer cell functionality and other immune responses.^{26,27}

The mortality rate among our IPA cases was 44.4%, and this changed to 41.3% after the cases from the literature were added. The mortality rate in critically ill patients with

TABLE 3 Organ support needs and complications between the two groups

	IPA (<i>n</i> = 63)	Non-IPA (<i>n</i> = 46)	<i>P</i> Value
<i>Organ support needs</i>			
IPPV, number (%)	45 (71.4)	34 (73.9)	0.774
IPPV days, days, median (IQR)	22 (9-27)	17 (7-18)	0.047*
Rescue ventilation strategies, ^a number (%)	27 (42.9)	10 (21.7)	0.021*
ECMO, number (%)	20 (31.7)	11 (23.9)	0.371
ECMO days, days, median (IQR)	11 (5-19)	13 (10-26)	0.455
Renal replacement therapy, number (%)	31 (49.2)	13 (28.3)	0.033*
Renal replacement therapy days, days, median (IQR)	7 (5-10)	7 (3-13)	0.677
High-dose vasoactive drugs, ^b number (%)	43 (68.3)	19 (41.3)	0.005*
<i>Complications during therapy</i>			
VAP, number (%)	32 (50.8)	20 (43.5)	0.450
CRBSI, number (%)	1 (1.6)	1 (2.2)	0.822
Bacteremia, number (%)	1 (1.6)	4 (8.7)	0.080
Barotrauma, number (%)	7 (11.1)	7 (15.2)	0.527
Haemorrhage, number (%)	7 (11.1)	6 (13.0)	0.759
Septic shock, number (%)	42 (66.7)	19 (41.3)	<0.001[#]

Abbreviations: CRBSI, catheter-related bloodstream infection; ECMO, extracorporeal membrane oxygenation; IPPV, invasive positive pressure ventilation; IQR, interquartile range; VAP, ventilator-associated pneumonia.

The bold values indicates *P* values from 0.01 to 0.05.

^aRescue ventilation strategies included the recruitment manoeuvre (RM), prone position ventilation (PP) and high-frequency oscillatory ventilation (HFOV).

^bHigh-dose vasoactive drugs refer to: noradrenaline >0.5 µg/kg/min or dopamine >20 µg/kg/min.

**P* < 0.05;

[#]*P* < 0.01.

influenza-associated IPA was reported to be 59%-95%.²⁸ A lack of awareness, the interference of underlying influenza, and the limited use of serological diagnostic methods in some medical institutions impeded the prompt diagnosis of IPA co-infection, delayed therapies and thus led to a high mortality rate. With the high incidence and mortality rates observed, awareness should be raised regarding influenza as an important risk factor for IPA co-infection, and medical care providers should watch for IPA during the course of influenza.

A multivariate logistic regression analysis suggested that the influenza patients who were treated with steroids after ICU admission, those with WBCs of more than $10 \times 10^9/L$ on ICU admission, and those whose CT imaging manifested as multiple nodules and cavities might have a high risk of IPA co-infection. According to reports, steroids are used in the treatment of 18%-69% of patients with influenza-associated acute respiratory distress syndrome (ARDS).²⁹ The benefits of limited inflammation in lung parenchyma with steroid use were not clear; however, the adverse effects, especially the immunosuppressive effects, might enable the growth of *Aspergillus* spp and hinder the efficacy of antifungal therapy.⁷ A study by Joost Wauters and his colleague⁷ revealed that the use of steroid 7 days before ICU admission is an independent risk factor for fungal co-infection. Currently, it

is well established that systemic corticosteroids increase the risk of IPA. In our study, the number of patients in the IPA group who received systemic steroids after ICU admission was 46%, which was much higher than in the non-IPA group, and the accumulated dose was up to 234 (40-400) mg, which was also higher than the level in the control group. Patients with influenza who are on corticosteroids before and/or after ICU admission should be carefully evaluated for possible *Aspergillus* co-infection; we also suggest that systemic steroids should be avoided in patients with influenza, even those with influenza-associated severe ARDS.

Patients with IPA in non-classic immunosuppressive hosts usually had high WBCs, which were distributed to the excessive inflammatory responses.²¹ This could account for the findings of multiple organ failure, such as acute kidney injury (49.2% vs 48.3, *P* = 0.033), and septic shock (66.7% vs 41.3, *P* < 0.001) that were observed more often in IPA patients compared with the control group. A WBC $>10 \times 10^9/L$ on ICU admission was an independent factor for IPA co-infection, especially when the procalcitonin (PCT) level, which indicates a bacterial infection, was normal. A lower lymphocyte count on ICU admission [$(0.47 \pm 0.52) \times 10^9/L$ vs $(0.67 \pm 0.46) \times 10^9/L$, *P* = 0.035], even if it is not an independent risk factor for IPA, should cause awareness of this

TABLE 4 Comparisons between survivors and non-survivors of IPA

	Survivors (<i>n</i> = 37)	Non-survivors (<i>n</i> = 26)	<i>P</i> Value
<i>General information</i>			
Age, years, mean ± SD	60 ± 15	67 ± 9	0.040*
Sex (men), number (%)	26 (70.3)	19 (73.1)	0.808
BMI > 30 kg/m ² , number (%)	2 (5.4)	3 (11.5)	0.375
SOFA, mean ± SD	8 ± 4	16 ± 4	<0.001#
Diabetes, number (%)	14 (37.8)	5 (19.2)	0.113
Underlying lung disease, number (%)	7 (18.9)	6 (23.1)	0.688
Chronic heart failure, number (%)	5 (13.5)	4 (15.4)	0.834
Chronic kidney disease, number (%)	2 (5.4)	0 (0.0)	0.228
Malignancy, number (%)	3 (8.1)	3 (11.5)	0.648
Immunosuppressant use in past 3 months, number (%)	10 (27.0)	5 (23.1)	0.474
<i>Systemic steroid use</i>			
Before ICU admission, number (%)	14 (37.8)	9 (34.6)	0.794
Accumulated dosage before ICU, mg, median (IQR)	391 (50-450)	892 (50-2365)	0.064
After ICU admission, number (%)	16 (43.2)	13 (50.0)	0.596
Accumulated dosage after ICU, mg, median (IQR)	509 (350-500)	574 (375-500)	0.520
Time from onset to IPA diagnosis, days, mean ± SD	9 (4-14)	10 (5-16)	0.518
<i>Laboratory examinations</i>			
Lymphocytes on ICU admission, *10 ⁹ /L, median (IQR)	0.54 (0.22-0.54)	0.38 (0.15-0.53)	0.271
CD4 ⁺ T cells on ICU admission, cell/μL, median (IQR)	315 (83-466)	152 (50-220)	0.031*
Influenza A, number (%)	26 (70.3)	26 (100.0)	0.002*
<i>Organ failure and support</i>			
IPPV, number (%)	20 (54.1)	25 (96.2)	<0.001#
Rescue ventilation strategies, number (%)	12 (32.4)	15 (57.7)	0.046*
ECMO, number (%)	7 (18.9)	13 (50.0)	0.015*
Renal replacement therapy, number (%)	17(45.9)	14 (53.8)	0.537
High-dose vasoactive drugs, number (%)	17 (45.9)	16 (61.5)	0.222

Abbreviations: BMI, body mass index; ECMO, extracorporeal membrane oxygenation; ICU, intensive care unit; IPPV, invasive positive pressure ventilation; IQR, interquartile range; SD, standard deviation; SOFA, sequential organ failure assessment; WBC, white blood cell.

The bold values indicates *P* values from 0.01 to 0.05.

Note. The steroid doses were converted to prednisone doses.

**P* < 0.05;

#*P* < 0.01.

possibility. Lymphopenia could lead to an alteration in the Th1 and Th2 balance, which was inclined to develop IPA.³

Radiological features in critically ill IPA patients are usually nonspecific, especially in patients with influenza-associated ARDS or bacterial co-infection. Our study suggested that multiple nodules were independent risk factors pointing to IPA in influenza patients. Multiple nodules distributed along the airway were common in non-classic immunosuppressive hosts.³⁰ These lung injuries are mainly due to an excessive host response rather than the fungal invasion itself, which mostly invades the vasculature, inducing secondary infraction; histological examinations of non-classic immunosuppressive patients often show large foci of pneumonia and exudative bronchiolitis with bronchial and alveolar

destruction. Thus, multiple nodules distributed along the airway were also relatively specific signs of IPA co-infection in influenza patients and should be recognized promptly.

Our study revealed that lower CD4⁺ T cell counts might predict a poor prognosis for IPA, which could be explained as patients lower CD4⁺ T cell counts might have weaker cellular immunity, making them much more vulnerable to *Aspergillus*.

Our study had some limitations. Its retrospective nature was the largest problem, along with some missing data might influence the expected statistical outcome in some degree. Although we included cases from after 2009, there was still heterogeneity among influenza strains, treatment strategies and geographic factors. Meanwhile, it is very difficult to suspect co-infections

TABLE 5 Multivariate logistic regression analysis for risk factors and predictors of prognosis for IPA in patients with influenza

	Wald	P value	Odds ratio	95% confidence interval	
				Inferior	Superior
<i>Risk factor for IPA co-infection</i>					
WBCs ($>10 \times 10^9/L$) on ICU admission	3.749	0.024	1.226	1.028	1.463
Steroids after ICU admission	3.568	0.045	5.372	0.980	29.460
CT indicated multiple nodules	11.902	0.001	9.833	1.965	49.194
CT indicated cavities	6.197	0.013	9.013	1.596	50.890
<i>Predictors of prognosis for IPA</i>					
SOFA	6.974	0.008	19.883	2.154	177.361
ECMO needed	5.211	0.022	1.210	1.027	1.432
CD4 ⁺ T cells (<200 cell/ μ L) on ICU admission	4.362	0.038	8.661	1.214	68.334

Abbreviations: CT, computed tomography; ICU, intensive care unit; SOFA, sequential organ failure assessment; WBC, white blood count.

and secondary infections in clinical practice; however, retrospective nature means data that might help identify features of co-infection was not consistently captured. Prospective, multi-centre studies are desired in order to elucidate the risk factors, clinical presentations and predictors of outcomes.

5 | CONCLUSIONS

With its high incidence and mortality, there should be greater awareness of influenza as an important risk factor for IPA co-infection. Influenza patients who receive steroids after ICU admission, who have WBCs of greater than $10 \times 10^9/L$ on ICU admission, and whose CT imaging shows multiple nodules and cavities might have a high risk of developing IPA. Additionally, poor prognoses may be predicted for patients with higher SOFA scores, CD4⁺ T cell counts lower than 200 cells/ μ L on ICU admission and more ECMO requirement.

AVAILABILITY OF DATA AND MATERIALS

The data sets used and/or analysed during the current study are available from the corresponding author upon reasonable request.

CONFLICT OF INTERESTS

The authors declare that they have no competing interests.

AUTHORS CONTRIBUTIONS

All authors made substantial contributions to the conception and design of the study or to the data acquisition, analysis or

interpretation; reviewed and approved the final manuscript; and significantly contributed to this study.

Study design: Qingyuan Zhan

Data collection: Linna Huang and Nannan Zhang

Drafting of the manuscript: Linna Huang and Nannan Zhang

Data verification and analysis: Linna Huang, Nannan Zhang, Xu Huang, Shuyu Xiong, Yingying Feng, Yi Zhang and Min Li.

ETHICS

This study was approved by the ethics committee of the China-Japan Friendship Hospital, Beijing, PR China and written informed consent was obtained from the patients or their next of kin.

ORCID

Linna Huang  <https://orcid.org/0000-0002-5682-8814>

REFERENCES

1. Alshabani K, Haq A, Miyakawa R, et al. Invasive pulmonary aspergillosis in patients with influenza infection: report of two cases and systematic review of the literature. *Expert Rev Respir Med.* 2015;9(1):89-96.
2. Brundage JF. Interactions between influenza and bacterial respiratory pathogens: implications for pandemic preparedness. *Lancet Infect Dis.* 2006;6(5):303-312.
3. Crum-Cianflone NF. Invasive aspergillosis associated with severe influenza infections. *Open Forum Infect Dis.* 2016;3(3):ofw171.
4. Shi SJ, Li H, Liu M, et al. Mortality prediction to hospitalized patients with influenza pneumonia: PO₂/FiO₂ combined lymphocyte count is the answer. *Clin Respir J.* 2017;11(3):352-360.

5. Lee JY, Joo EJ, Yeom JS, et al. Aspergillus tracheobronchitis and influenza A co-infection in a patient with AIDS and neutropenia. *Infect Chemother*. 2014;46(3):209-215.
6. Lat A, Bhadelia N, Miko B, et al. Invasive aspergillosis after pandemic (H1N1) 2009. *Emerg Infect Dis*. 2010;16(6):971-973.
7. Wauters J, Baar I, Meersseman P, et al. Invasive pulmonary aspergillosis is a frequent complication of critically ill H1N1 patients: a retrospective study. *Intensive Care Med*. 2012;38(11):1761-1768.
8. Kwon OK, Lee MG, Kim HS, et al. Invasive pulmonary aspergillosis after influenza a infection in an immunocompetent patient. *Tuberc Respir Dis (Seoul)*. 2013;75(6):260-263.
9. Toh HS, Jiang MY, Tay HT. Invasive pulmonary aspergillosis in severe complicated influenza A. *J Formos Med Assoc*. 2013;112(12):810-811.
10. Otterspoor LC, Smit FH, van Laar TJ, et al. Prolonged use of extracorporeal membrane oxygenation combined with prone positioning in patients with acute respiratory distress syndrome and invasive aspergillosis. *Perfusion*. 2012;27(4):335-337.
11. Vehreschild JJ, Brockelmann PJ, Bangard C, et al. Pandemic 2009 influenza A(H1N1) virus infection coinciding with invasive pulmonary aspergillosis in neutropenic patients. *Epidemiol Infect*. 2012;140(10):1848-1852.
12. Hoyo-Ulloa I, Cobos-Trigueros N, Puig-de la Bellacasa J, et al. Influenza A (H1N1) complicated by invasive aspergillosis in non-severely immunocompromised patients. *Enferm Infecc Microbiol Clin*. 2012;30(9):583-584.
13. Guervilly C, Roch A, Ranque S, et al. A strategy based on galactomannan antigen detection and PCR for invasive pulmonary aspergillosis following influenza A (H1N1) pneumonia. *J Infect*. 2012;65(5):470-473.
14. Adalja AA, Sappington PL, Harris SP, et al. Isolation of Aspergillus in three 2009 H1N1 influenza patients. *Influenza Other Respir Viruses*. 2011;5(4):225-229.
15. Kim SH, Kim MN, Lee SO, et al. Fatal pandemic influenza A/H1N1 infection complicated by probable invasive pulmonary aspergillosis. *Mycoses*. 2012;55(2):189-192.
16. Garcia-Vidal C, Barba P, Arnan M, et al. Invasive aspergillosis complicating pandemic influenza A (H1N1) infection in severely immunocompromised patients. *Clin Infect Dis*. 2011;53(6):e16-19.
17. Carfagna P, Brandimarte F, Caccese R, et al. Occurrence of influenza A(H1N1)v infection and concomitant invasive pulmonary aspergillosis in a patient with chronic obstructive pulmonary disease. *Mycoses*. 2011;54(6):549-551.
18. Passouant O, Mateu P, Commandini M, et al. Pulmonary aspergillosis in non-immunocompromised patient with acute respiratory distress syndrome during A (H1N1) infection. *Ann Fr Anesth Reanim*. 2011;30(11):e75-76.
19. Gao R, Cao B, Hu Y, et al. Human infection with novel avian-origin influenza A (H7N9) virus. *N Engl J Med*. 2013;368(20):1888-1897.
20. De Pauw B, Walsh TJ, Donnelly JP, et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis*. 2008;46(12):1813-1821.
21. Bulpa P, Dive A, Sibille Y. Invasive pulmonary aspergillosis in patients with chronic obstructive pulmonary disease. *Eur Respir J*. 2007;30(4):782-800.
22. Blot SI, Taccone FS, Van den Abeele, et al. A clinical algorithm to diagnose invasive pulmonary aspergillosis in critically ill patients. *Am J Respir Crit Care Med*. 2012;186(1):56-64.
23. Huang L, He H, Zhan Q, et al. Is Bulpa criteria suitable for the diagnosis of probable invasive pulmonary Aspergillosis in critically ill patients with chronic obstructive pulmonary disease? A comparative study with EORTC/MSG and ICU criteria. *BMC Infect Dis*. 2017;17(1):209.
24. Guarner J, Falcon-Escobedo R. Comparison of the pathology caused by H1N1, H5N1, and H3N2 influenza viruses. *Arch Med Res*. 2009;40(8):655-661.
25. Writing Committee of the WHO CoCAoPI. Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *N Engl J Med*. 2010;362(18):1708-1719.
26. Lee N, Wong CK, Chan PK, et al. Cytokine response patterns in severe pandemic 2009 H1N1 and seasonal influenza among hospitalized adults. *PLoS ONE*. 2011;6(10):e26050.
27. Fox A, Le NM, Horby P, et al. Severe pandemic H1N1 2009 infection is associated with transient NK and T deficiency and aberrant CD8 responses. *PLoS ONE*. 2012;7(2):e31535.
28. Dimopoulos G, Frantzeskaki F, Poulakou G, et al. Invasive aspergillosis in the intensive care unit. *Ann N Y Acad Sci*. 2012;1272:31-39.
29. Brun-Buisson C, Richard JC, Mercat A, et al. Early corticosteroids in severe influenza A/H1N1 pneumonia and acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2011;183(9):1200-1206.
30. Huang L, He H, Zhan Q, et al. Values of radiological examinations for the diagnosis and prognosis of invasive bronchial-pulmonary aspergillosis in critically ill patients with chronic obstructive pulmonary diseases. *Clin Respir J*. 2018;12(2):499-509.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

FILE S1. Modified Bulpa Criteria. Probable IPA in our study was diagnosed using the 'Modified Bulpa Criteria'.²⁴

How to cite this article: Huang L, Zhang N, Huang X, et al. Invasive pulmonary aspergillosis in patients with influenza infection: A retrospective study and review of the literature. *Clin Respir J*. 2019;13: 202–211. <https://doi.org/10.1111/crj.12995>